

Secant varieties of Segre-Veronese varieties

Dustin Cartwright ¹

October 6, 2011

¹joint with Daniel Erman and Luke Oeding

Tensor rank for matrices

U, V : finite dimensional vector spaces

$$x \in U \otimes V$$

The **rank** of x is the smallest integer r such that x can be written

$$x = u_1 \otimes v_1 + \cdots + u_r \otimes v_r \text{ where } u_i \in U, v_i \in V.$$

Tensor rank for matrices

U, V : finite dimensional vector spaces

$$x \in U \otimes V$$

The **rank** of x is the smallest integer r such that x can be written

$$x = u_1 \otimes v_1 + \cdots + u_r \otimes v_r \text{ where } u_i \in U, v_i \in V.$$

- ▶ The rank can **jump down** but never up at special points.

Tensor rank for matrices

U, V : finite dimensional vector spaces

$$x \in U \otimes V$$

The **rank** of x is the smallest integer r such that x can be written

$$x = u_1 \otimes v_1 + \cdots + u_r \otimes v_r \text{ where } u_i \in U, v_i \in V.$$

- ▶ The rank can **jump down** but never up at special points.
- ▶ The rank is the same if we pass to a **bigger field**.

Tensor rank for matrices

U, V : finite dimensional vector spaces

$$x \in U \otimes V$$

The **rank** of x is the smallest integer r such that x can be written

$$x = u_1 \otimes v_1 + \cdots + u_r \otimes v_r \text{ where } u_i \in U, v_i \in V.$$

- ▶ The rank can **jump down** but never up at special points.
- ▶ The rank is the same if we pass to a **bigger field**.
- ▶ The set of **possible decompositions** is a homogeneous space.

Partially symmetric tensors

U : m -dimensional \mathbb{C} -vector space

V : n -dimensional \mathbb{C} -vector space

$$x \in U \otimes S^2V$$

The **rank** of x is the smallest integer r such that x can be written in the form:

$$x = u_1 \otimes v_1 \otimes v_1 + \cdots + u_r \otimes v_r \otimes v_r$$

Partially symmetric tensors

U : m -dimensional \mathbb{C} -vector space

V : n -dimensional \mathbb{C} -vector space

$$x \in U \otimes S^2 V$$

The **rank** of x is the smallest integer r such that x can be written in the form:

$$x = u_1 \otimes v_1 \otimes v_1 + \cdots + u_r \otimes v_r \otimes v_r$$

The rank can jump both down and **up** for special tensors.

Partially symmetric tensors

U : m -dimensional \mathbb{C} -vector space

V : n -dimensional \mathbb{C} -vector space

$$x \in U \otimes S^2 V$$

The **border rank** of x is the smallest integer r such that x can be approximated arbitrarily closely by expressions of the form:

$$x \approx u_1 \otimes v_1 \otimes v_1 + \cdots + u_r \otimes v_r \otimes v_r$$

Partially symmetric tensors

U : m -dimensional \mathbb{C} -vector space

V : n -dimensional \mathbb{C} -vector space

$$x \in U \otimes S^2 V$$

The **border rank** of x is the smallest integer r such that x can be approximated arbitrarily closely by expressions of the form:

$$x \approx u_1 \otimes v_1 \otimes v_1 + \cdots + u_r \otimes v_r \otimes v_r$$

The set of such decompositions will in general be a finite set of points, possibly defined over a **larger field** than x .

Equations for bounded border rank

When $U = \mathbb{C}e_1 \oplus \mathbb{C}e_2$, we can write

$$x = e_1 \otimes A + e_2 \otimes B$$

where A and B are symmetric matrices.

Equations for bounded border rank

When $U = \mathbb{C}e_1 \oplus \mathbb{C}e_2$, we can write

$$x = e_1 \otimes A + e_2 \otimes B$$

where A and B are symmetric matrices.

Theorem

*The ideal of partially symmetric tensors whose border rank is at most r is generated by the $(r + 1) \times (r + 1)$ -minors of the **block matrix***

$$\begin{pmatrix} A & B \end{pmatrix}$$

Equations for small border rank

When $U = \mathbb{C}e_1 \oplus \mathbb{C}e_2 \oplus \mathbb{C}e_3$, we can write

$$x = e_1 \otimes A + e_2 \otimes B + e_3 \otimes C$$

Equations for small border rank

When $U = \mathbb{C}e_1 \oplus \mathbb{C}e_2 \oplus \mathbb{C}e_3$, we can write

$$x = e_1 \otimes A + e_2 \otimes B + e_3 \otimes C$$

Theorem (C.-Erman-Oeding 2010)

If $r \leq 5$, the ideal of tensors whose border rank is at most r is generated by the $(r+1) \times (r+1)$ -*minors* and $(2r+2) \times (2r+2)$ -*Pfaffians* respectively of

$$(A \quad B \quad C) \quad \text{and} \quad \begin{pmatrix} 0 & A & -B \\ -A & 0 & C \\ B & -C & 0 \end{pmatrix}$$

Equations for small border rank

When $U = \mathbb{C}e_1 \oplus \mathbb{C}e_2 \oplus \mathbb{C}e_3$, we can write

$$x = e_1 \otimes A + e_2 \otimes B + e_3 \otimes C$$

Theorem (C.-Erman-Oeding 2010)

If $r \leq 5$, the ideal of tensors whose border rank is at most r is generated by the $(r+1) \times (r+1)$ -*minors* and $(2r+2) \times (2r+2)$ -*Pfaffians* respectively of

$$(A \quad B \quad C) \quad \text{and} \quad \begin{pmatrix} 0 & A & -B \\ -A & 0 & C \\ B & -C & 0 \end{pmatrix}$$

Remark

The $n = 4$, $r = 5$ case is due to *Emil Toeplitz* in 1869.

Outline of the proof

- ▶ Assume $n = r$ and A is the identity matrix. Then

$$\begin{pmatrix} 0 & I & -B \\ -I & 0 & C \\ B & -C & 0 \end{pmatrix} \sim \begin{pmatrix} 0 & I & 0 \\ -I & 0 & 0 \\ 0 & 0 & BC - CB \end{pmatrix}$$

The $2r + 2$ -Pfaffians of this matrix are the entries of the commutator $BC - CB$, which is a prime, Gorenstein ideal, defining the variety of **commuting symmetric matrices**.

Outline of the proof

- ▶ Assume $n = r$ and A is the identity matrix. Then

$$\begin{pmatrix} 0 & I & -B \\ -I & 0 & C \\ B & -C & 0 \end{pmatrix} \sim \begin{pmatrix} 0 & I & 0 \\ -I & 0 & 0 \\ 0 & 0 & BC - CB \end{pmatrix}$$

The $2r + 2$ -Pfaffians of this matrix are the entries of the commutator $BC - CB$, which is a prime, Gorenstein ideal, defining the variety of **commuting symmetric matrices**.

- ▶ Now just assume $n = r$. We need to **bound the dimension** of the set of tensors where A is singular. This is computational and is only true for $r \leq 5$.

Outline of the proof

- ▶ Assume $n = r$ and A is the identity matrix. Then

$$\begin{pmatrix} 0 & I & -B \\ -I & 0 & C \\ B & -C & 0 \end{pmatrix} \sim \begin{pmatrix} 0 & I & 0 \\ -I & 0 & 0 \\ 0 & 0 & BC - CB \end{pmatrix}$$

The $2r + 2$ -Pfaffians of this matrix are the entries of the commutator $BC - CB$, which is a prime, Gorenstein ideal, defining the variety of **commuting symmetric matrices**.

- ▶ Now just assume $n = r$. We need to **bound the dimension** of the set of tensors where A is singular. This is computational and is only true for $r \leq 5$.
- ▶ Arbitrary n . Here the minors come in.

Unifying framework for these equations

Given decomposable $u \otimes v \otimes v \in U \otimes S^2V$, we have linear map

$$\begin{aligned}\psi_{j,u \otimes v \otimes v}: V^* \otimes \bigwedge^j U &\rightarrow V \otimes \bigwedge^{j+1} U \\ v^* \otimes (u'_1 \wedge \cdots \wedge u'_j) &\mapsto \langle v^*, v \rangle v \otimes u'_1 \wedge \cdots \wedge u'_j \wedge u\end{aligned}$$

Unifying framework for these equations

Given decomposable $u \otimes v \otimes v \in U \otimes S^2V$, we have linear map

$$\begin{aligned}\psi_{j,u \otimes v \otimes v}: V^* \otimes \bigwedge^j U &\rightarrow V \otimes \bigwedge^{j+1} U \\ v^* \otimes (u'_1 \wedge \cdots \wedge u'_j) &\mapsto \langle v^*, v \rangle v \otimes u'_1 \wedge \cdots \wedge u'_j \wedge u\end{aligned}$$

For arbitrary $x \in U \otimes S^2V$, define $\psi_{j,x}$ by **extending linearly**.

Unifying framework for these equations

Given decomposable $u \otimes v \otimes v \in U \otimes S^2V$, we have linear map

$$\begin{aligned}\psi_{j,u \otimes v \otimes v}: V^* \otimes \bigwedge^j U &\rightarrow V \otimes \bigwedge^{j+1} U \\ v^* \otimes (u'_1 \wedge \cdots \wedge u'_j) &\mapsto \langle v^*, v \rangle v \otimes u'_1 \wedge \cdots \wedge u'_j \wedge u\end{aligned}$$

For arbitrary $x \in U \otimes S^2V$, define $\psi_{j,x}$ by **extending linearly**.
If U is 3-dimensional,

- ▶ The $j = 0$ and $j = 2$ cases give the **rectangular matrix**
- ▶ The $j = 1$ case gives the **skew-symmetric** square matrix

Robust testing of determinantal equations

Let

$$\sigma_1 \geq \cdots \geq \sigma_4 \quad \text{and} \quad \sigma'_1 \geq \cdots \geq \sigma'_{12}$$

be the **singular values** of

$$(A \ B \ C) \quad \text{and} \quad \begin{pmatrix} 0 & A & -B \\ -A & 0 & C \\ B & -C & 0 \end{pmatrix}$$

respectively.

Robust testing of determinantal equations

Let

$$\sigma_1 \geq \cdots \geq \sigma_4 \quad \text{and} \quad \sigma'_1 \geq \cdots \geq \sigma'_{12}$$

be the **singular values** of

$$(A \quad B \quad C) \quad \text{and} \quad \begin{pmatrix} 0 & A & -B \\ -A & 0 & C \\ B & -C & 0 \end{pmatrix}$$

respectively.

The functions

$$\gamma_r^2 = \sum_{i=r+1}^4 \sigma_i^2 \qquad \delta_r^2 = \sum_{i=2r+1}^{12} (\sigma'_i)^2$$

Robust testing of determinantal equations

Let

$$\sigma_1 \geq \cdots \geq \sigma_4 \quad \text{and} \quad \sigma'_1 \geq \cdots \geq \sigma'_{12}$$

be the **singular values** of

$$(A \ B \ C) \quad \text{and} \quad \begin{pmatrix} 0 & A & -B \\ -A & 0 & C \\ B & -C & 0 \end{pmatrix}$$

respectively.


The functions

$$\gamma_r^2 = \sum_{i=r+1}^4 \sigma_i^2 \qquad \delta_r^2 = \sum_{i=2r+1}^{12} (\sigma'_i)^2$$

are **continuous**, non-negative functions which are both zero if and only if the tensor has rank at most r .


Bounded real rank is a semi-algebraic set

Tensors with **real** border rank at most r characterized by same equalities, but additional **inequalities**



Bounded real rank is a semi-algebraic set

Equalities are more important than inequalities for detecting deviations



Thank you