

Self-Intersections of Closed Geodesics on a
Negatively Curved Surface: Statistical Regularities

by

Steven P. Lalley
Department of Statistics
Purdue University

Technical Report # 92-08

Department of Statistics
Purdue University

February 1992

SELF-INTERSECTIONS OF CLOSED GEODESICS ON A
NEGATIVELY CURVED SURFACE: STATISTICAL REGULARITIES

by

Steven P. Lalley
Department of Statistics
Purdue University

ABSTRACT

For compact surfaces of negative curvature it is shown that most closed geodesics of length $\approx \ell$ have about $C\ell^2$ self-intersections, for some constant $C > 0$, and these self-intersections are approximately equidistributed on the surface. For surfaces of constant negative curvature $-\kappa$ the value of the constant C is $\kappa/2\pi^2(g-1)$, where g is the genus.

1. Introduction and Statement of Main Results

The geodesic flow on a compact, negatively curved surface is perhaps the simplest example of a smooth flow for which typical orbits exhibit “random” behavior. Periodic orbits (i.e., closed geodesics) are *not* typical – there are only countably many, and they account for only a set of measure zero in the phase space. Nevertheless, the overall randomness of the flow is reflected in the sequence of periodic orbits (see, e.g., [L₁], [L₂]) in certain ways. The purpose of this note is to record some statistical regularities in the self-intersections of closed geodesics.

Let S be a compact, C^∞ Riemannian manifold of dimension 2 with strictly negative curvature at every point. There are countably many closed geodesics $\gamma_1, \gamma_2, \dots$ (one in each free homotopy class) with lengths $\ell_1 \leq \ell_2 \leq \dots$. A celebrated result of Margulis [M] states that if $\pi(t) = \max\{n: \ell_n \leq t\}$ then as $t \rightarrow \infty$

$$\pi(t) \sim \frac{e^{ht}}{ht}$$

where $h > 0$ is the topological entropy of the geodesic flow (for surfaces with constant curvature -1 , $h = 1$).

Define s_n to be the number of (transversal) self-intersections of the closed geodesic γ_n .

Theorem 1: *For each compact, negatively curved surface S there is a constant $C = C_S > 0$ such that for every $\varepsilon > 0$,*

$$(1.1) \quad \lim_{t \rightarrow \infty} \pi(t)^{-1} |\{n \leq \pi(t): |s_n - C\ell_n^2| < \varepsilon\ell_n^2\}| = 1$$

If S has constant curvature $-\kappa$ and genus $g \geq 2$ then

$$(1.2) \quad C_S = \frac{\kappa}{2\pi^2(g-1)}.$$

Thus, “most” closed geodesics of length $\approx \ell$ have “about” $C_S \ell^2$ self-intersections. This result is consistent with the statement that “typical” closed geodesics are similar statistically to “generic” geodesics. Observe that if one randomly threw down ℓ geodesic segments of length 1 on S then the number of intersections would be about $C\ell^2$.

Theorem 1 seems to contradict Theorem 5 of [P], which states that most closed geodesics of length $\approx \ell$ have about $\tilde{C}\ell$ self-intersections. Apparently, the formula in remark (iii), p. 212 of [P] is incorrect: the singularity in the zeta function is not a simple pole, as stated, but rather a logarithmic singularity, so the Ikehara Tauberian theorem does *not* apply.

If $s_n > 0$ define α_n to be the probability distribution on S that assigns mass $\frac{1}{s_n}$ to each point of self-intersection of γ_n ; if $s_n = 0$ define α_n to be the zero measure.

Theorem 2: For each compact, negatively curved surface S there is a Borel probability measure α on S such that for each continuous $f: S \rightarrow \mathbf{R}$ and $\varepsilon > 0$

$$(1.3) \quad \lim_{t \rightarrow \infty} \pi(t)^{-1} \sum_{n=1}^{\pi(t)} 1\{|\int f d\alpha_n - \int f d\alpha| < \varepsilon\} = 1.$$

If S has constant curvature then $\alpha = \nu =$ normalized area measure on S .

Thus most closed geodesics on S of length about ℓ experience about $C_S \ell^2$ self-intersections and these self-intersections are approximately distributed according to α . In fact the proof will show that α is the projection to S of the maximum entropy invariant probability measure for the geodesic flow. Theorem 1 is proved in sec. 3, Th. 2 in sec. 4. Both theorems are in fact corollaries of the strong equidistribution result, Theorem 7, of [L₁]. This result is explained in sec. 2.

Simple closed geodesics (closed geodesics with no self-intersections) are of interest for both topological and number-theoretic reasons (see [S]). Theorem 2 shows that simple closed geodesics are atypical of closed geodesics. Results of Birman and Series [BS] suggest that at least for a noncompact surface with finite area and free fundamental group, the number of simple closed geodesics with period $\leq t$ grows no faster than polynomially in t ; thus simple closed geodesics are *very* atypical. It would be interesting to have an asymptotic formula for simple closed geodesics analogous to Margulis' formula.

2. Equidistribution of Closed Geodesics

We begin by discussing Theorem 7 of [L₁], from which Theorems 1-2 will be deduced.

Let $A = A(i, j)$ be an irreducible, aperiodic $\ell \times \ell$ matrix of zeros and ones, where $\ell \geq 2$, and define

$$\Sigma_A = \{x \in \prod_{n=-\infty}^{\infty} \{1, 2, \dots, \ell\} : A(x_n, x_{n+1}) = 1 \ \forall n\}.$$

The shift $\sigma: \Sigma_A \rightarrow \Sigma_A$ is defined by $(\sigma x)_n = x_{n+1}$. Let $r: \Sigma_A \rightarrow \mathbf{R}$ be a strictly positive function which is Lipschitz relative to the metric d_ρ on Σ_A defined by

$$d_\rho(x, y) = \sum_{n=-\infty}^{\infty} 1\{x_n \neq y_n\} \rho^{|n|}$$

for some $\rho \in (0, 1)$. Define the suspension space

$$\Sigma_A^r = \{(x, s) : x \in \Sigma_A \text{ and } 0 \leq s \leq r(x)\}$$

with the points $(x, r(x))$ and $(\sigma x, 0)$ identified. The *suspension flow* (Σ_A^r, σ_t^r) , $-\infty < t < \infty$, is defined as follows. Starting at any $(x, s) \in \Sigma_A^r$, move at unit speed up the fiber

$(x, s'), s \leq s' \leq r(x)$, until reaching $(x, r(x))$, then jump instantaneously to $(\sigma x, 0)$ and proceed up the vertical fiber $(\sigma x, s')$, etc. Equivalently,

$$\begin{aligned}\sigma_t^r(x, s) &= (x, s + t) \quad \forall 0 \leq s \leq s + t \leq r(x), \\ \sigma_{t+t'}^r &= \sigma_t^r \circ \sigma_{t'}^r.\end{aligned}$$

Observe that the periodic orbits of the suspension flow (Σ_A^r, σ^r) are precisely those orbits which pass through some $(x, 0)$ with x a periodic sequence. Since there are countably many periodic sequences, there are countably many periodic orbits of (Σ_A^r, σ^r) . These may be labelled $\gamma_1^*, \gamma_2^*, \dots$ and their periods $\ell_1^* \leq \ell_2^* \leq \dots$. The distribution ν_n^* of γ_n^* may be defined by

$$\nu_n^*(B) = \int_0^{\ell_n^*} 1_B(\gamma_n^*(t)) dt$$

where $B \subset \Sigma_A^r$ is a Borel set. Set $\pi^*(t) = \max\{n: \ell_n \leq t\}$.

Theorem 7 ([L₁]): As $t \rightarrow \infty$,

$$\pi^*(t) \sim e^{ht}/ht$$

where h is the topological entropy of (Σ_A^r, σ^r) . For each continuous $f: \Sigma_A^r \rightarrow \mathbf{R}$ and each $\varepsilon > 0$, as $t \rightarrow \infty$

$$\pi^*(t)^{-1} |\{n \leq \pi^*(t): |\int f d\nu_n^* - \int f d\nu^*| < \varepsilon\}| \rightarrow 1,$$

where ν^* is the maximum entropy invariant measure for (Σ_A^r, σ^r) .

Now consider the geodesic flow $\Phi = \Phi_t$ on the unit tangent bundle T_1S of a compact, negatively curved surface. This flow is of Anosov type [A], hence also Axiom A, so the results of [B₂] (also [R]) apply. Thus, there exists a suspension flow (Σ_A^r, σ^r) and a continuous map $\pi: \Sigma_A^r \rightarrow T_1S$ such that

- (a) π is surjective;
- (b) π is at most N to 1 for some $N < \infty$;
- (c) $\pi \circ \sigma_t^r = \Phi_t \circ \pi$ for all t ;
- (d) all but finitely many of the periodic orbits $\{\bar{\gamma}_n\}$ of Φ have the property that $\pi^{-1}(\bar{\gamma}_n)$ consists of a single periodic orbit of σ^r with the same least period.

See [L₃], sec. 1 for an explanation of (d). Note that the periodic orbits $\bar{\gamma}_n$ of Φ are just the lifts to T_1S of the closed geodesics γ_n . For each $\bar{\gamma}_n$ the probability distribution $\bar{\nu}_n$ may be defined by

$$\int_{T_1S} f d\bar{\nu}_n = \frac{1}{\ell_n} \int_0^{\ell_n} f(\bar{\gamma}_n(t)) dt$$

for $f: T_1S \rightarrow \mathbf{R}$ continuous. By (a)–(d) above, $\bar{\nu}_n$ pulls back (via π^{-1}) to $\nu_{m_n}^*$, with only finitely many exceptions. Furthermore, the maximum entropy invariant probability

measure $\bar{\nu}$ for the geodesic flow Φ pulls back to the maximum entropy measure ν^* for the suspension flow σ^r . Therefore, Theorem 1 implies

Corollary 1: *For each continuous $f: T_1S \rightarrow \mathbf{R}$ and each $\varepsilon > 0$,*

$$\lim_{t \rightarrow \infty} \frac{1}{\pi(t)} |\{n \leq \pi(t) : |\int f d\bar{\nu}_n - \int f d\bar{\nu}| < \varepsilon\}| = 1.$$

3. The Number of Self-Intersections

Recall that $\bar{\nu}$ is the maximum entropy invariant probability measure for the geodesic flow on T_1S , and that ν is the induced measure on S .

Lemma 1: *For any nonempty, open set $U \subset T_1S$, $\bar{\nu}(U) > 0$. For any orbit $\bar{\gamma}(t)$ of the flow, if $\Gamma = \{\bar{\gamma}(t) : -\infty < t < \infty\}$ then $\bar{\nu}(\Gamma) = 0$.*

Note: This is part of the folklore, but the proof is not easy to find.

Proof: It suffices to prove corresponding statements for an arbitrary suspension flow, by (a)–(c) of sec. 2. Let ν^* be the maximum entropy measure for a suspension flow (Σ_A^r, σ^r) ; define a probability measure μ on Σ_A by

$$\mu(F) = \nu^*\{(x, s) : x \in F \text{ and } 0 \leq s \leq r(x)\}.$$

Then μ is the equilibrium state (sometimes called Gibbs state) for the function $-hr(x)$, by [BR], Prop. 3 (see [B₃], Ch. 1 for the definition).

We will prove that any Gibbs state μ is nonatomic. From this it follows immediately that ν^* assigns measure zero to each individual orbit of the flow. Suppose μ has an atom x , i.e., $\mu(\{x\}) = \rho > 0$. Since μ is σ -invariant ([B₁], Th. (1.2)), $\mu(\{\sigma^n x\}) = \rho \ \forall n$; as μ is a finite measure, it must be that $x = \sigma^n x$ for some $n \geq 1$. But μ is mixing ([B₁], Prop. (1.14)) so it must be that $\mu(\{x\}) = 1$ and $x = \sigma x$. This is impossible, however, because by [B₁] Th. (1.2) a Gibbs state μ gives positive mass to every cylinder set.

It remains to show that $\nu^*(W) > 0$ for every open $W \subset \Sigma_A^r$. If W is open then it contains a rectangle $R = F \times [k/m, (k+1)/m)$, where k, m are positive integers and F is a cylinder set $F = \{y \in \Sigma_A : y_n = x_n \ \forall |n| \leq n_*\}$. By [B₁] Th. (1.2), $\mu(F) > 0$. Since ν^* is σ_t^r -invariant,

$$\nu^*(F \times [0/m, 1/m)) = \nu^*(F \times [1/m, 2/m)) = \dots$$

(with the obvious convention about what happens when you get to the “ceiling” $\{(x, r(x))\}$: see Fig. 1). Let k_* be the smallest integer larger than $\min_{x \in F} r(x)$; then

$$\mu(F) \leq \sum_{i=0}^{k_*-1} \nu^*(F \times [i/m, (i+1)/m)) - k_* \nu_*(R)$$

(see Fig. 1 again) so $\nu^*(R) > 0$. □

Figure 1

Lemma 2: *For any geodesic $\gamma(t)$, if $G = \{\gamma(t): -\infty < t < \infty\}$ then $\nu(G) = 0$.*

Proof: The maximum entropy measure $\bar{\nu}$ is ergodic for the geodesic flow on T_1S (this follows from (a)–(c) of sec. 2, because the maximum entropy measure for a suspension flow is unique and consequently ergodic). Hence, if $\nu(G) > 0$ then for $\bar{\nu}$ -a.e. orbit $\bar{\varphi}(t) = (\varphi(t), \varphi'(t))$ of the geodesic flow

$$\lim_{T \rightarrow \infty} T^{-1} \int_0^T 1_G(\varphi(t)) dt = \nu(G) > 0.$$

Now $\varphi(t)$ has at most countably many transversal intersections with G , and each transversal intersection contributes zero to the integral $\int_0^T 1_G(\varphi(t)) dt$. Therefore $\varphi(t)$ must intersect G tangentially. But $\varphi(t)$ and $\gamma(t)$ are both geodesics, so it follows that for some $s \in \mathbf{R}$, $\varphi(t) = \gamma(t+s) \forall t$. This is impossible, because by Lemma 1 $\bar{\nu}$ assigns zero mass to the orbit $\{\bar{\gamma}(t) = (\gamma(t), \gamma'(t))\}$. □

Each $(x, v) \in T_1S$ determines a geodesic $\gamma(t)$ in S emanating from $\gamma(0) = x$ and with $\gamma'(0) = v$. For any $\delta > 0$ let $G_\delta(x, v)$ be the segment $\{\gamma(t): 0 \leq t < \delta\}$ of this geodesic, considered as a subset of S . Let $\bar{\nu} \times \bar{\nu}$ denote the product measure on $T_1S \times T_1S$ determined by $\bar{\nu} \times \bar{\nu}(B_1 \times B_2) = \bar{\nu}(B_1)\bar{\nu}(B_2)$. Define

$$U_\delta = \{((x, v), (x', v')): G_\delta(x, v) \cap G_\delta(x', v') \neq \emptyset\};$$

then $\bar{\nu} \times \bar{\nu}(U_\delta)$ is the probability that two independent, randomly chosen geodesic segments of length δ cross. (Note that the $\bar{\nu} \times \bar{\nu}$ -probability of a nontransversal intersection is zero, by Lemma 1.)

Lemma 3: For each $\delta > 0$, $\bar{\nu} \times \bar{\nu}(U_\delta) > 0$.

Proof: Each $U_\delta, \delta > 0$, contains a nonempty open subset of $T_1S \times T_1S$, consequently also a rectangle $A \times B$ where A, B are nonempty open subsets of T_1S . Therefore, by Lemma 1, U_δ has positive $\bar{\nu} \times \bar{\nu}$ -measure. \square

When $\bar{\mu}$ is close to $\bar{\nu}$ in the weak-* topology then $\bar{\mu} \times \bar{\mu}$ is close to $\bar{\nu} \times \bar{\nu}$ in the weak-* topology, and consequently $|\bar{\mu} \times \bar{\mu}(U_\delta) - \bar{\nu} \times \bar{\nu}(U_\delta)|$ should be small. The following lemma justifies this assertion (cf. [Bi], Th. 2.1, statement (v)).

Lemma 4: Let ∂U_δ denote the (topological) boundary of U_δ in $T_1S \times T_1S$. Then

$$\bar{\nu} \times \bar{\nu}(\partial U_\delta) = 0.$$

Proof: If $((x, v), (x', v')) \in \partial U_\delta$ then an endpoint of $G_\delta(x, v)$ lies on $G_\delta(x', v')$, or an endpoint of $G_\delta(x', v')$ lies on $G_\delta(x, v)$. Consequently, to prove the lemma it suffices to show that for each $(x', v') \in T_1S$,

$$\bar{\nu}\{(x, v): G_\delta(x, v) \text{ has an endpoint on } G_\delta(x', v')\} = 0.$$

For $(x, v) \in T_1S$, one endpoint of $G_\delta(x, v)$ is x ; call the other x_* . By Lemma 2, $\bar{\nu}\{(x, v): x \in G_\delta(x', v')\} = 0$. But $\bar{\nu}$ is invariant under time reversal, and the segment $G_\delta(x, v)$ reversed has initial endpoint x_* , so $\bar{\nu}\{(x, v): x_* \in G_\delta(x, v)\} = 0$. \square

Geodesics, being smooth curves, look like straight lines in the small. Therefore, if $\delta > 0$ is sufficiently small then two geodesic segments $G_\delta(x, v)$ and $G_\delta(x', v')$ of length δ will have at most one transverse intersection (this also uses the compactness of T_1S). Choose δ_0 sufficiently small that this is true for all $0 < \delta < \delta_0$. For the remainder of this section, fix δ with $0 < \delta < \delta_0$.

Let γ_n be a closed geodesic with length ℓ_n . Recall that s_n is the number of (transversal) self-intersections of γ_n , and that $\bar{\nu}_n$ is the distribution of $\bar{\gamma}_n = (\gamma_n, \gamma'_n)$ in T_1S . For $k = 1, 2, \dots$ let $m_k (= m_k^{(n)})$ be the least integer $\geq k\ell_n/\delta$. Define

$$\begin{aligned} x_{j,k} &= x_{j,k}^{(n)} = \gamma_n(j\delta/k), j = 0, 1, 2, \dots, m_k - 1; \\ v_{j,k} &= v_{j,k}^{(n)} = \gamma'_n(j\delta/k), j = 0, 1, 2, \dots, m_k - 1; \\ \bar{\nu}_{n,k} &= \text{uniform probability distribution on}\{(x_{j,k}, v_{j,k}): j = 0, 1, \dots, m_k - 1\} \end{aligned}$$

(i.e., $\bar{\nu}_{n,k}$ is the probability measure that puts mass $1/m_k$ on each $x_{j,k}, v_{j,k}, j = 0, 1, \dots, m_k - 1$).

Lemma 5: weak-* $\lim_{k \rightarrow \infty} \bar{\nu}_{n,k} = \bar{\nu}_n$.

This is immediate from the definition of the weak-* topology and elementary properties of the Riemann integral.

Lemma 6: For each $k = 1, 2, \dots$,

$$\begin{aligned} & (m_k^2/2k^2)(\bar{\nu}_{n,k} \times \bar{\nu}_{n,k})(U_\delta) - 2m_k/k \leq \\ & s_n \leq (m_k^2/2k^2)(\bar{\nu}_{n,k} \times \bar{\nu}_{n,k})(U_\delta). \end{aligned}$$

Proof: Define G_j to be the segment of γ_n from $x_{j,k}$ to $x_{j+1,k}$ for $j = 0, 1, \dots, m_k - 2$, and G_{m_k-1} to be the segment of γ_n from $x_{m_k-1,k}$ to $x_{0,k}$. Note that if $0 \leq j \leq m_k - 2$ then $G_j = G_{\delta/k}(x_{j,k}, v_{j,k})$, whereas $G_{m_k-1} = G_r(x_{m_k-1,k}, v_{m_k-1,k})$ for some $0 < r \leq \delta/k$. Since $\delta < \delta_0$, each intersection $G_i \cap G_j$, $i \neq j$, consists of at most a single point. Consequently,

$$s_n = \frac{1}{2} \sum_{(i,j): i \neq j} 1\{G_i \cap G_j \neq \emptyset\}.$$

Next, define $G_i^* = G_\delta(x_{i,k}, v_{i,k})$. If $0 \leq i < m_k - k$ then $G_i^* = G_i \cup G_{i+1} \cup \dots \cup G_{i+k-1}$, whereas if $m_k - k \leq i \leq m_k - 1$ then $G_i^* \supset G_i \cup G_{i+1} \cup \dots \cup G_{i+k-1}$ but $G_i^* \subset G_i \cup G_{i+1} \cup \dots \cup G_{i+k}$ (with the convention that if $i + j \geq m_k$ then $i + j$ should be reduced mod m_k). As before, each intersection $G_i^* \cap G_j^*$, $|i - j| \geq k$, consists of at most one point. Each G_i is contained in k distinct G_j^* , so by the result of the previous paragraph,

$$\begin{aligned} s_n & \leq \frac{1}{2k^2} \sum_{(i,j): |i-j| \geq k} 1\{G_i^* \cap G_j^* \neq \emptyset\} \\ & \leq \frac{m_k^2}{2k^2} (\bar{\nu}_{n,k} \times \bar{\nu}_{n,k})(U_\delta). \end{aligned}$$

Moreover, each G_i intersects only k distinct G_j^* unless $0 \leq i < k$, in which case G_i may intersect $k + 1$ distinct G_j^* . Consequently,

$$\left(\frac{1}{2k^2} \sum_{(i,j): |i-j| \geq k} 1\{G_i^* \cap G_j^* \neq \emptyset\} \right) - s_n \leq \frac{2km_k}{2k^2} = \frac{m_k}{k}.$$

Finally,

$$\begin{aligned} & \frac{m_k^2}{2k^2} (\bar{\nu}_{n,k} \times \bar{\nu}_{n,k})(U_\delta) - \frac{1}{2k^2} \sum_{(i,j): |i-j| \geq k} 1\{G_i^* \cap G_j^* \neq \emptyset\} \\ & = \frac{1}{2k^2} \sum_{i=0}^{m_k-1} (2k-1) = \frac{(2k-1)m_k}{2k^2} \leq \frac{m_k}{k}. \quad \square \end{aligned}$$

Proof of (1.1): Corollary 1 implies that for most closed geodesics γ_n with length $\ell_n \leq t$ the distribution $\bar{\nu}_n$ is close to $\bar{\nu}$ in the weak-* topology. Consequently, by Lemma 5, for k sufficiently large $\bar{\nu}_{n,k}$ is close to $\bar{\nu}$ and therefore $\bar{\nu}_{n,k} \times \bar{\nu}_{n,k}$ is close to $\bar{\nu} \times \bar{\nu}$ in the weak-*

topology. It follows, by Lemma 4 and [Bi], Th. 2.1, that $|\bar{\nu}_{n,k} \times \bar{\nu}_{n,k}(U_\delta) - \bar{\nu} \times \bar{\nu}(U_\delta)|$ is small. Thus, by Lemma 6, s_n/ℓ_n^2 is close to

$$C = \bar{\nu} \times \bar{\nu}(U_\delta)/\delta^2 > 0. \quad \square$$

Since $\delta < \delta_0$ was arbitrary in this argument, the quantity $\bar{\nu} \times \bar{\nu}(U_\delta)/\delta^2$ is independent of δ , and so C can, in principle, be evaluated by letting $\delta \rightarrow 0$.

Lemma 7: *If S has constant negative curvature then*

$$(3.1) \quad \lim_{\delta \rightarrow 0} \bar{\nu} \times \bar{\nu}(U_\delta)/\delta^2 = \frac{4}{2\pi \text{ area}(S)}$$

Observe that $\text{area}(S) = 4\pi(g-1)/\kappa$ where $g = \text{genus}(S)$ and $-\kappa = \text{curvature}$, by the Gauss-Bonnet theorem, so Lemma 7 implies (1.2).

Proof of Lemma 7: The measure $\bar{\nu}$ has a simple structure for a surface S of constant curvature. Elements of $T_1 S$ may be represented as (x, θ) , where $x \in S$ and $-\pi \leq \theta < \pi$; in these coordinates

$$d\bar{\nu}(x, \theta) = d\nu(x)d\theta/2\pi$$

where $d\nu(x)$ is the normalized surface area measure on S .

Recall that $\bar{\nu} \times \bar{\nu}(U_\delta)$ is the probability that two independent, randomly chosen geodesic segments of length δ will intersect. To prove (3.1) it suffices to show that for any fixed geodesic segment G_δ of length δ

$$\bar{\nu}\{(x, \theta): G_\delta(x, \theta) \cap G_\delta \neq \emptyset\} \sim 4\delta^2/2\pi \text{ area}(S)$$

as $\delta \rightarrow 0$ (uniformly for all choices of G_δ). To compute this probability, condition on the value of θ (the angle with G_δ); then the set of x for which $G_\delta(x, \theta) \cap G_\delta \neq \emptyset$ is approximately a rhombus of side δ and interior angle θ (at least when δ is small). Consequently,

$$\begin{aligned} & \bar{\nu}\{(x, \theta): G_\delta(x, \theta) \cap G_\delta \neq \emptyset\} \\ & \sim \int_{-\pi}^{\pi} \delta^2 |\sin \theta| d\theta / 2\pi \text{ area}(S). \quad \square \end{aligned}$$

4. Distribution of Self-Intersection Points

In sec. 3 we showed that $\bar{\nu} \times \bar{\nu}(U_\delta) = C\delta^2$ for all sufficiently small $\delta > 0$. Thus we may define Borel probability measures μ_δ on $T_1 S \times T_1 S$ by

$$\mu_\delta(B) = \bar{\nu} \times \bar{\nu}(U_\delta \cap B)/C\delta^2$$

for Borel sets $B \subset T_1 S$. Since $T_1 S \times T_1 S$ is compact, the Helly selection theorem (cf. [RS], Th. IV.21, which they call the ‘‘Banach-Alaoglu’’ theorem) implies that there is a weak-* convergent subsequence $\mu_{\delta_n}, \delta_n \rightarrow 0$. Define

$$\bar{\alpha} = \text{weak-}^* \lim_{n \rightarrow \infty} \mu_{\delta_n}$$

(it will not matter which subsequence δ_n is used).

The measure $\bar{\alpha}$ induces on $S \times S$ a measure α via the natural projection $p \times p: T_1 S \rightarrow S \times S$, in particular,

$$\alpha(B) = \bar{\alpha}((p \times p)^{-1}(B)), B \subset S \text{ Borel.}$$

The measure α is supported by the ‘‘diagonal’’ $D = \{(x, x): x \in S\}$, because μ_{δ} is supported by the set U_{δ} , and if $((x, v), (x', v')) \in U_{\delta}$ then distance $(x, x') \leq 2\delta$. Thus α may be regarded as a probability measure on S . We will not bother to distinguish between S and D in the subsequent arguments.

Let V be an open subset of $S \times S$; define

$$V_{\delta}^+ = \{(x_1, x_2) \in S \times S: \text{distance}(x_i, V) < \delta\},$$

$$V_{\delta}^- = \{(x_1, x_2) \in V: \text{distance}(x_i, V^c) > \delta\}.$$

Lemma 8: *If $\alpha(\partial V) = 0$ then*

$$\lim_{n \rightarrow \infty} \mu_{\delta_n}((p \times p)^{-1}(V_{2\delta_n}^+)) = \alpha(V) \text{ and}$$

$$\lim_{n \rightarrow \infty} \mu_{\delta_n}((p \times p)^{-1}(V_{\delta_n}^-)) = \alpha(V).$$

bf Proof: This is a standard argument. For any $\varepsilon > 0$ there exist open $V_1 \subset V \subset V_2$ such that $\alpha(V_2 \setminus V_1) < \varepsilon$ and $V_1 \subset V_{\delta}^-, V_2 \supset V_{\delta}^+$ for some $\delta > 0$ (this follows from the dominated convergence theorem), and also such that $\alpha(\partial V_2) = \alpha(\partial V_1) = 0$. Since $\mu_{\delta_n} \rightarrow \bar{\alpha}$ it then follows ([Bi], Th. 2.1 (v)) that

$$\mu_{\delta_n}((p \times p)^{-1}(V_1)) \rightarrow \alpha(V_1),$$

$$\mu_{\delta_n}((p \times p)^{-1}(V_2)) \rightarrow \alpha(V_2).$$

The result then follows by monotonicity, because for large n , $V_1 \subset V_{\delta_n}^- \subset V \subset V_{2\delta_n}^+ \subset V_2$. \square

Now let γ_n be a closed geodesic with length ℓ_n , and let $\bar{v}_{n,k}, m_k$, etc. be as in Lemma 6. If V is any closed subset of $S \times S$ let $s_n(V)$ be the number of self-intersection points x of γ_n such that $(x, x) \in V$.

Lemma 9: *For each $k = 1, 2, \dots$,*

$$\begin{aligned} & (m_k^2/2k^2)(\bar{v}_{n,k} \times \bar{v}_{n,k})(U_{\delta} \cap (p \times p)^{-1}(V_{\delta}^-)) - 2m_k/k \\ & \leq s_n(V) \leq (m_k^2/2k^2)(\bar{v}_{n,k} \times \bar{v}_{n,k})(U_{\delta} \cap (p \times p)^{-1}(V_{\delta}^+)). \end{aligned}$$

Proof: This is virtually the same as the proof of Lemma 6. The only novelty is in keeping track of where the geodesic segments cross. Observe that if geodesic segments G_1, G_2 of length δ intersect transversally at x such that $(x, x) \in V$ then the (initial) endpoints x_1, x_2 of G_1, G_2 are such that $(x_1, x_2) \in V_\delta^+$. Similarly, if G_1, G_2 intersection at x and the initial endpoints x_1, x_2 are such that $(x_1, x_2) \in V_\delta^-$, then $(x, x) \in V$. \square

Proof of Th. 2: Let V be any open subset of $S \times S$ such that $\alpha(\partial V) = 0$. By Th. 1, Lemma 5, and Lemma 9, for any $\varepsilon > 0$ there is a $t_\varepsilon < \infty$ large enough that for all $t \geq t_\varepsilon$ and all but at most $\varepsilon\pi(t)$ closed geodesics γ_n with length $\leq t$,

$$C\mu_\delta((p \times p)^{-1}(V_\delta^-)) - \varepsilon \leq s_n(V)/\ell_n^2 \leq C\mu_\delta((p \times p)^{-1}(V_\delta^+)) + \varepsilon.$$

The quantities ε and δ are both arbitrary, but affect how large t must be so that the preceding statement is true. By letting $\delta \rightarrow 0$ through the subsequence δ_n and appealing to Lemma 8, we see that for any $\varepsilon > 0$ there exists $t_\varepsilon < \infty$ such that for all $t \geq t_\varepsilon$ and all but at most $\varepsilon\pi(t)$ closed geodesics γ_n of length $\leq t$,

$$C\alpha(V) - 2\varepsilon \leq s_n(V)/\ell_n^2 \leq C\alpha(V) + 2\varepsilon.$$

Together with Th. 1, this implies that for most closed geodesics γ_n with $\ell_n \leq t$ the distribution α_n of self-intersection points is close to α in the weak-* topology. \square

REFERENCES

- [A] Anosov, D.V. (1967). Geodesic flows on closed Riemannian manifolds of negative curvature. *Proc. Steklov Inst.* **90**.
- [Bi] Billingsley, P. (1968). *Convergence of Probability Measures*. Wiley, N.Y.
- [BS] Birman, J. and Series, C. (1984). An algorithm for simple curves on surfaces. *J. London Math. Soc.* (2) **29**, 331–342.
- [B₁] Bowen, R. (1975). *Equilibrium States and the Ergodic Theorem of Anosov Diffeomorphisms*. Springer Lecture Notes in Math. **470**
- [B₂] Bowen, R. (1973). Symbolic dynamics for hyperbolic flows. *Amer. J. Math.* **95**, 429–450.
- [BR] Bowen, R. and Ruelle, D. (1975). Ergodic theory of Axiom A flows. *Inv. Math.* **29**, 181–202.
- [L₁] Lalley, S.P. (1989). Renewal theorems in symbolic dynamics, with applications to geodesic flows, noneuclidean tessellations and their fractal limits. *Acta Mathematica* **163**, 1–55.
- [L₂] Lalley, S.P. (1987). Distribution of periodic orbits of symbolic and Axiom A flows. *Adv. Appl. Math.* **8**, 154–193.
- [L₃] Lalley, S.P. (1989). Closed geodesics in homology classes on surfaces of variable negative curvature. *Duke Math. J.* **58**, 795–821.
- [M] Margulis, G. (1969). Applications of ergodic theory to the investigation of manifolds of negative curvature. *Funct. Analysis Appl.* **3**, 335–336.
- [P] Pollicott, M. (1985). Asymptotic distribution of closed geodesics. *Israel J. Math.* **52**, 209–224.
- [R] Ratner, M. (1973). Markov partitions for Anosov flows on n -dimensional manifolds. *Israel J. Math.* **15**, 92–114.
- [RS] Reed, M., and Simon, B. (1980). *Methods of Mathematical Physics I. Functional Analysis*. Academic Press, N.Y.
- [S] Series, C. (1985). The geometry of Markov numbers. *Math. Intell.* **7**, 20–29.